

Nur für Forschungszwecke

# Caspase 9/p35/p10 Polyklonaler Antikörper

Katalog-Nr.: CL647-10380



## Allgemeine Informationen

Katalog-Nr.:	CL647-10380	GenBank-Zugangsnummer:	BC002452
Größe:	100ul , Konzentration: 1000 µg/ml von 842 Nanodrop;	GenID (NCBI):	Anregungs-/Emissionsmaxima-Wellenlängen:
Wirz:	Kaninchen	Vollständiger Name:	654 nm / 674 nm
Isotyp:	IgG	Berechneté Masse:	46 kDa
Immunogen Katalognummer:	AG0404	Beobachteté Masse:	46 kDa, 35 kDa

## Anwendungen

Geprüfte Anwendungen:  
FC (Intra)

Getestete Reaktivität:  
Human, Maus, Ratte

## Hintergrundinformationen

Caspase 9, apoptosis-related cysteine protease (CASP9, synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c) is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Caspase 9 is processed by APAF1; this step is thought to be one of the earliest in the caspase activation cascade. 10380-1-AP can recognize the pre- and cleaved- caspase 9. In recent years, the localization of caspase9 was a focus of interest. Beside its cytoplasmic distribution, a very extensive localization study was done on rat brain tissue, where caspase9 was found located predominantly in the nucleus and to a lesser extend in the cytoplasm [PMID: 15541731].

## Lagerung


Lagerungsbedingungen:  
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.  
Lagerungspuffer:  
BS mit 50% Glyzerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.  
Aliquotieren ist nicht notwendig bei -20°C Lagerung

\*\*\* 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Ausgewählte Validierungsdaten



$1 \times 10^6$  HepG2 cells were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human Caspase 9/p35/p10 (CL647-10380) (red), or 0.2 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).