

Nur für Forschungszwecke

NF-κB p65 Polyklonaler Antikörper

Katalog-Nr.: CL647-10745

Vorgestelltes Produkt



Allgemeine Informationen

Katalog-Nr.:	CL647-10745	GenBank-Zugangsnummer:	BC011603
Größe:	100ul , Konzentration: 1000 µg/ml von 5970 Nanodrop;	GenID (NCBI):	
Wirz:	Kaninchen	Vollständiger Name:	v-rel reticuloendotheliosis viral oncogene homolog A (avian)
Isotyp:	IgG	Berechneté Masse:	65 kDa
Immunogen Katalognummer:	AG1199	Beobachteté Masse:	65 kDa

Reinigungsmethode:
Antigen-Affinitätsreinigung
Anregungs-/Emissionsmaxima-Wellenlängen:
654 nm / 674 nm

Anwendungen

Geprüfte Anwendungen:
FC (Intra)

Getestete Reaktivität:
Human, Maus, Ratte

Hintergrundinformationen

Nuclear factor κB (NF-κB) is a sequence-specific DNA-binding protein complex which regulates the expression of viral genomes, including the human immunodeficiency virus, and a variety of cellular genes, particularly those involved in immune and inflammatory responses. The members of the NF-κB family in mammalian cells include the proto-oncogene c-Rel, p50/p105 (NFKB1), p65 (RelA), p52/p100 (NFKB2), and RelB. All of these proteins share a conserved 300-amino acid region known as the Rel homology domain which is responsible for DNA binding, dimerization, and nuclear translocation of NF-κB. The p65 subunit is a major component of NF-κB complexes and is responsible for trans-activation. NF-κB heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-κB p65-p65 complex appears to be involved in invasin-mediated activation of IL-8 expression. The inhibitory effect of IκB upon NF-κB in the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-κB complex. It associates with chromatin at the NF-κB promoter region via association with DDX1. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human RELA.

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.
Lagerungspuffer:
BS mit 50% Glyzerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human NF-κB p65 (CL647-10745) (red), or 0.2 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).