

Nur für Forschungszwecke

HDAC1 Monoklonaler Antikörper

Katalog-Nr.:CL647-66085



Allgemeine Informationen

Katalog-Nr.: CL647-66085	GenBank-Zugangsnummer: BC000301	Reinigungsmethode: Protein-G-Reinigung
Größe: 100ul , Konzentration: 1000 µg/ml von3065 Nanodrop;	GeneID (NCBI): 3065	CloneNo.: 4C4G7
Wirt: Maus	Vollständiger Name: histone deacetylase 1	Empfohlene Verdünnungen: IF 1:500-1:2000
Isotyp: IgG1	Berechnete Masse: 55 kDa	Anregungs-/Emissionsmaxima- Wellenlängen: 654 nm / 674 nm
Immunogen Katalognummer: AG19128	Beobachtete Masse: 65 kDa	

Anwendungen

Geprüfte Anwendungen: FC (Intra), IF	Positivkontrollen: IF : A431-Zellen,
Getestete Reaktivität: Human, Maus, Ratte	

Hintergrundinformationen

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC 1 is 55 kDa, but modified HDAC 1 is about 60-65 kDa .

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Vor Licht schützen.
Lagerungspuffer:
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite® Plus 647 HDAC1 antibody (CL647-66085, Clone: 4C4G7) at dilution of 1:1000.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 647 Anti-Human HDAC1 (CL647-66085, Clone:4C4G7) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).