

Nur für Forschungszwecke

# TNF Alpha Monoklonaler Antikörper

Katalog-Nr.:CL647-69002



## Allgemeine Informationen

<b>Katalog-Nr.:</b> CL647-69002	<b>GenBank-Zugangsnummer:</b> GeneID (NCBI): 7124	<b>Reinigungsmethode:</b> Protein-G-Reinigung
<b>Größe:</b> 100ul , Konzentration: 1000 µg/ml von Nanodrop;	<b>Vollständiger Name:</b> tumor necrosis factor (TNF superfamily, member 2)	<b>CloneNo.:</b> 11H8A5
<b>Wirt:</b> Maus		<b>Anregungs-/Emissionsmaxima- Wellenlängen:</b> 654 nm / 674 nm
<b>Isotyp:</b> IgG1		
<b>Immunogen Katalognummer:</b> HZ-1014		

## Anwendungen

**Geprüfte Anwendungen:**  
FC (Intra)

**Getestete Reaktivität:**  
Human

## Hintergrundinformationen

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, INS resistance, and cancer.

## Lagerung

**Lagerungsbedingungen:**  
Bei -20°C lagern. Vor Licht schützen. Nach dem Versand ein Jahr stabil.

**Lagerungspuffer:**  
BS mit 50% Glycerin, 0,05% Proclin300, 0,5% BSA, pH 7,3.

**Aliquotieren ist nicht notwendig bei -20°C Lagerung**

**\*\*\* 20ul-Größen enthalten 0.1% BSA**




For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Ausgewählte Validierungsdaten



1X10<sup>6</sup> PMA+ionomycin-stimulated and Monensin Solution treated human PBMCs were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5), and 0.25 ug CoraLite®488 Anti-Human CD3 (UCHT1) (CL488-65151, Clone: UCHT1). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

1X10<sup>6</sup> human PBMCs were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5), and 0.25 ug CoraLite®488 Anti-Human CD3 (UCHT1) (CL488-65151, Clone: UCHT1). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

1X10<sup>6</sup> NK92 were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5) (red), or 0.25 ug x. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).