Nur für Forschungszwecke

CD64 Monoklonaler Antikörper

Katalog-Nr.:CR-65253



Allgemeine Informationen

Katalog-Nr.:

CR-65253 Größe:

Wirt:

100tests , 5 μ l/test

Maus Isotyp:

IgG1, kappa

Berechneté Masse: 374 aa, 43 kDa

receptor (CD64)

BC032634

2209

GeneID (NCBI):

Vollständiger Name:

GenBank-Zugangsnummer:

Fc fragment of IgG, high affinity Ia,

Reinigungsmethode:

Affinitätsreinigung CloneNo.:

Anregungs-/Emissionsmaxima-

Wellenlängen: 592 nm / 611 nm

Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Human

Hintergrundinformationen

Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Lagerung


Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Lagerungspuffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Ausgewählte Validierungsdaten



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD64 (CR-65253, Clone:10.1) (red) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD64 (CR-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.