Nur für Forschungszwecke

CD11c Monoklonaler Antikörper

Katalog-Nr.:FITC-65086



Reinigungsmethode:

Anregungs-/Emissionsmaxima-

Affinitätsreinigung

CloneNo.:

Wellenlängen: 495 nm / 524 nm

Allgemeine Informationen

Katalog-Nr.:

FITC-65086 Größe:

100 tests , 5 μ l/test

Wirt: Maus

Isotyp: IgG1, kappa

Geprüfte Anwendungen:

Getestete Reaktivität:

Human

Hintergrundinformationen

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α/β heterodimer with CD18 (integrin β2). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

GenBank-Zugangsnummer:

integrin, alpha X (complement

component 3 receptor 4 subunit)

BC038237

3687

GeneID (NCBI):

Vollständiger Name:

Berechneté Masse: 1169 aa, 129 kDa

Lagerung

Anwendungen


Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Lagerungspuffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Ausgewählte Validierungsdaten



1X10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD11c (FITC-65086, Clone: 3.9) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD11c (FITC-65086, Clone: 3.9) or unstained. Cells were not fixed. Monocytes were gated.