

Nur für Forschungszwecke

CD107b / LAMP2 Monoklonaler Antikörper

Katalog-Nr.: PE-65053



Allgemeine Informationen

Katalog-Nr.: PE-65053	GenBank-Zugangsnummer: BC002965	Reinigungsmethode: Affinitätsreinigung
Größe: 100tests , 10 µl/test	GeneID (NCBI): 3920	CloneNo.: H4B4
Wirt: Maus	Vollständiger Name: lysosomal-associated membrane protein 2	Empfohlene Verdünnungen: IF 1:50-1:500
Isotyp: IgG1, kappa	Berechnete Masse: 45 kDa	Anregungs-/Emissionsmaxima-Wellenlängen: 496 nm, 565 nm / 578 nm

Anwendungen

Geprüfte Anwendungen: FC (Intra), IF	Positivkontrollen: IF : HeLa-Zellen,
Getestete Reaktivität: Human	

Hintergrundinformationen

LAMP2 (CD107b) is a Lysosomal membrane glycoprotein. LAMP2 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP-2 is localized primarily in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane under certain circumstances, e.g., after platelet activation, during granulocytic differentiation and activation, and in some tumor cells (PMID: 12221139). LAMP is involved in lysosomal stability and autophagy (PMID: 12221139). This glycoprotein provides selectins with carbohydrate ligands. LAMP2 may play a role in tumor cell metastasis (PMID 9426697).

Lagerung

Lagerungsbedingungen:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Lagerungspuffer:
PBS mit 0,1% Natriumazid und einem Stabilisator.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PE-65053 (CD107b antibody) at dilution of 1:50 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ Jurkat cells were intracellularly stained with 10 ug PE Anti-Human CD107b (PE-65053, Clone:H4B4) (red), or 10 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).