## Nur für Forschungszwecke

## CD28 Monoklonaler Antikörper

Katalog-Nr.:PE-65099



Allgemeine Informationen

Katalog-Nr.: PE-65099

Größe:

100 tests , 5  $\mu$ l/test

Wirt: Maus Isotyp:

IgG1, kappa

GenBank-Zugangsnummer:

BC093698 GeneID (NCBI):

940

Vollständiger Name: CD28 molecule Berechneté Masse:

220 aa, 25 kDa

Reinigungsmethode:

Affinitätsreinigung

CloneNo.: CD28.2

Anregungs-/Emissionsmaxima-

Wellenlängen:

496 nm, 565 nm / 578 nm

Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Human

Hintergrundinformationen

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).

Lagerung


Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Lagerungspuffer:

PBS with 0.09% sodium azide.

## Ausgewählte Validierungsdaten



1X10^6 human PBMCs were surface co-stained with CD3 and 5 ul PE Anti-Human CD28 (PE-65099, Clone:CD28.2) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD28 (PE-65099, Clone:CD28.2) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.