

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TRAIL



Numéro de catalogue: 10399-1-AP

1 Publications

Informations de base

Numéro de catalogue:
10399-1-AP

Taille:
150ul , Concentration: 220 µg/ml by
Bradford method using BSA as the
standard;

Hôte:
Lapin

Isotype:
IgG

Immunogen Catalog Number:
AG0627

Numéro d'acquisition GenBank:
BC032722

Identification du gène (NCBI):
8743

Nom complet:
tumor necrosis factor (ligand)
superfamily, member 10

MW calculé
281 aa, 33 kDa

Méthode de purification:
Purification par affinité contre
l'antigène

Dilutions recommandées:
WB 1:300-1:600

Applications

Applications testées:
WB, ELISA

Demandes citées:
WB

Spécificité de l'espèce:
Humain

Espèces citées:
Humain

Contrôles positifs:

WB : cellules K-562,

Informations générales

TRAIL is a 30 kD protein known as TNF-related apoptosis inducing ligand, CD253, TNFSF10, or Apo-2L. It is expressed as two kinds of forms: cell surface protein (30 kD) and soluble ligand (19-20 kD) produced by megakaryocytes and platelets, monocytes, neutrophils, NK cells and activated T cells. This antigen is also expressed on tumor cells and transformed cell lines. TRAIL has been reported to induce apoptosis in tumor and transformed cell lines by a caspase-dependent process.

Publications notables

Autrice	Pubmed ID	Journal	Application
Hui Li	30384810	Technol Cancer Res Treat	WB

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



K-562 cells were subjected to SDS PAGE followed by western blot with 10399-1-AP (TRAIL antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.