

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-ASC/TMS1



Numéro de catalogue: 10500-1-AP

Phare

178 Publications

Informations de base

Numéro de catalogue:

10500-1-AP

Taille:

150ul, Concentration: 600 µg/ml by Nanodrop;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG0774

Numéro d'acquisition GenBank:

BC004470

Identification du gène (NCBI):

29108

Nom complet:

PYD and CARD domain containing

MW calculé

25 kDa

MW observés:

22-25 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:5000-1:50000

IP 0.5-4.0 µg for IP and 1:2000-1:16000 for WB

IHC 1:50-1:500

Applications

Applications testées:

IHC, IP, WB, ELISA

Demandes citées:

CoIP, IF, IHC, IP, WB

Spécificité de l'espèce:

Humain

Espèces citées:

bovin, canin, Humain, porc

Contrôles positifs:

WB : cellules HL-60, cellules A549, cellules Raji, cellules THP-1, cellules U-937

IP : cellules HL-60,

IHC : tissu de cancer du poumon humain, tissu de côlon de souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Informations générales

PYCARD, also known as ASC or TMS1, is a 195 amino acid protein containing an N-terminal Pyrin-like domain (PYD) and an 87 residue C-terminal Caspase-associated recruitment domain (CARD). It promotes caspase-mediated apoptosis which is mediated predominantly through the activation of caspase-9.

Publications notables

Autrice	Pubmed ID	Journal	Application
JiangFan Yu	36248980	Front Oncol	WB
Sheng-Li Ming	33072119	Front Immunol	WB
Pengxiao Chen	34603026	Front Pharmacol	WB,IF

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 10500-1-AP (ASC/TMS1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-ASC/TMS1(IP:10500-1-AP, 4ug; Detection:10500-1-AP 1:8000) with HL-60 cells lysate 1720 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 10500-1-AP (TMS1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).