

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-RTN4/NOGO



Numéro de catalogue: 10740-1-AP

14 Publications

## Informations de base

Numéro de catalogue:

10740-1-AP

Taille:

150ul, Concentration: 500 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG1164

Numéro d'acquisition GenBank:

BC007109

Identification du gène (NCBI):

57142

Nom complet:

reticulon 4

MW calculé

130 kDa

MW observés:

190-210 kDa, 45 kDa, 22-25 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:8000

IHC 1:50-1:500

## Applications

Applications testées:

FC, IHC, WB, ELISA

Demandes citées:

IF, IP, WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

Humain, rat, souris

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) A défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

Contrôles positifs:

WB : tissu cérébral de souris, tissu cérébral de rat

IHC : tissu testiculaire de souris, tissu de cancer de la prostate humain

## Informations générales

Reticulon (RTN) proteins are a group of membrane-bound proteins that largely reside in endoplasmic reticulum (ER) (PMID: 18177508). Reticulon proteins share a common sequence feature, the reticulon homology domain (RHD). They are involved in shaping the tubular endoplasmic reticulum network, membrane trafficking, inhibition of axonal growth, and apoptosis (PMID: 24218324). Four mammalian reticulons (RTN1-4) exist. RTN4 (also known as Neurite outgrowth inhibitor or Nogo) is a myelin-associated neurite growth inhibitory protein. Some isoforms of RTN4 have been described. RTN4A (Nogo-A, runs at ~200 kDa), RTN4B (Nogo-B, 40-55 kDa), and RTN4C (Nogo-C, 22-25 kDa) are three major isoforms (PMID: 31092426; 16469703).

## Publications notables

Autrice	Pubmed ID	Journal	Application
Maria Sol Herrera-Cruz	34743744	Biol Direct	WB
Xiao Wang	33186557	Cell Metab	WB
Dwayne J Byrne	35346366	Acta Neuropathol Commun	WB,IF

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 10740-1-AP (RTN4/NOGO antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 10740-1-AP (RTN4/NOGO antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 10740-1-AP (RTN4/NOGO antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10<sup>6</sup> SH-SY5Y cells were stained with 0.2ug RTN4/NOGO antibody (10740-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.