

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-CD69



Numéro de catalogue: 10803-1-AP

5 Publications

Informations de base

Numéro de catalogue:	BC007037	Méthode de purification:
10803-1-AP		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
150ul , Concentration: 750 µg/ml by Nanodrop;	969	WB 1:500-1:3000
Hôte:	Nom complet:	IHC 1:2000-1:8000
Lapin	CD69 molecule	IF 1:50-1:500
Isotype:	MW calculé	
IgG	23 kDa	
Immunogen Catalog Number:	MW observés:	
AG1094	28 kDa, 32 kDa	

Applications

Applications testées:	Contrôles positifs:
IF, IHC, WB, ELISA	WB: plaquettes de sang périphérique humain,
Demandes citées:	IHC : tissu d'amygdalite humain,
FC, IF, IHC, WB	IF : tissu d'amygdalite humain,
Spécificité de l'espèce:	
Humain	
Espèces citées:	
Humain	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>	

Informations générales

CD69, also known as activation inducer molecule (AIM) or early activation antigen (EA1), is a cell surface glycoprotein expressed shortly after activation on T lymphocytes and other cells of haematopoietic origin, including natural killer cells. It is a disulfide-linked homodimer composed of two chains belonging to the type II integral protein with a C-type lectin domain. CD69 is involved in lymphocyte proliferation and functions as a signal-transmitting receptor in lymphocytes, natural killer cells, and platelets.

Publications notables

Autrice	Pubmed ID	Journal	Application
Shijia Rao	31661988	Expert Rev Clin Immunol	IF
Huiwen Zheng	35594870	Cell Rep	IF
Chen Zhu	31428519	Oncoimmunology	IHC

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 10803-1-AP (CD69 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD69 antibody (10803-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD69 antibody (10803-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).