

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-RTN4/NOGO



Numéro de catalogue: 10950-1-AP

Phare

8 Publications

Informations de base

Numéro de catalogue:	BC012619	Méthode de purification:
10950-1-AP		Purification par affinité contre l'antigène
Taille:	57142	Dilutions recommandées:
150ul , Concentration: 600 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	Nom complet: reticulon 4	WB 1:5000-1:50000
Hôte:	MW calculé	IHC 1:50-1:500
Lapin	130 kDa	IF 1:10-1:100
Isotype:	MW observés:	
IgG	45-50 kDa	
Immunogen Catalog Number:		
AG1392		

Applications

Applications testées:	Contrôles positifs:
FC, IF, IHC, WB, ELISA	WB : cellules A375, cellules A549, cellules HepG2, cellules SH-SY5Y
Demandes citées:	IHC : tissu cérébral de souris, tissu testiculaire humain
IF, WB	IF : cellules A375,
Spécificité de l'espèce:	
Humain, souris	
Espèces citées:	
Humain, rat, souris	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.</i>	

Informations générales

Reticulon (RTN) proteins are a group of membrane-bound proteins that largely reside in endoplasmic reticulum (ER) (PMID: 18177508). Reticulon proteins share a common sequence feature, the reticulon homology domain (RHD). They are involved in shaping the tubular endoplasmic reticulum network, membrane trafficking, inhibition of axonal growth, and apoptosis (PMID: 24218324). Four mammalian reticulons (RTN1-4) exist. RTN4 (also known as Neurite outgrowth inhibitor or Nogo) is a myelin-associated neurite growth inhibitory protein. Some isoforms of RTN4 have been described. RTN4A (Nogo-A, runs at ~200 kDa), RTN4B (Nogo-B, 40-55 kDa), and RTN4C (Nogo-C, 22-25 kDa) are three major isoforms (PMID: 31092426; 16469703).

Publications notables

Autrice	Pubmed ID	Journal	Application
Jing Wang	32764877	Drug Des Devel Ther	WB
Tamutenda Chidawanyika	33622772	J Cell Sci	WB
Nicholas J Lennemann	28102736	Autophagy	WB,IF

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 10950-1-AP (RTN4/NOGO antibody) at dilution of 1:120000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10950-1-AP (RTN4/NOGO antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of A375 cells using 10950-1-AP (RTN4/NOGO antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human RTN4/NOGO (10950-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 10950-1-AP (RTN4/NOGO antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).