

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TNFSF13B

Numéro de catalogue:[11862-1-AP](#)



Informations de base

Numéro de catalogue:	BC020674	Méthode de purification:
11862-1-AP		Purification par affinité contre l'antigène
Taille:	10673	Dilutions recommandées:
150ul , Concentration: 600 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	tumor necrosis factor (ligand) superfamily, member 13b	WB 1:500-1:2000
Hôte:	MW calculé	
Lapin	32 kDa	
Isotype:	MW observés:	
IgG	32 kDa, 45-52 kDa	
Immunogen Catalog Number:		
AG2440		

Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules U-937, tissu hépatique de rat, tissu hépatique de souris, tissu hépatique humain, tissu testiculaire de souris
Spécificité de l'espèce:	
Humain, rat, souris	

Informations générales

TNFSF13B, also known as BAFF, BLYS, TALL1, TNFSF20, ZTNF4 and CD257, is a cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. This cytokine is expressed in B lineage cells, and acts as a potent B cell activator. The BAFF system promotes B cell survival and differentiation and plays a prominent role in the pathogenesis of autoimmune diseases. It has been shown that BAFF inhibitors are in clinical trials for systemic lupus erythematosus with significant efficacy. This antibody recognizes all the forms of TNFSF13B such as soluble form (17 kDa), membrane form (32-35 kDa) and glycosylation form (45-52 kDa).

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 11862-1-AP (TNFSF13B antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



U-937 cells were subjected to SDS PAGE followed by western blot with 11862-1-AP (TNFSF13B antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.