

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-DOK5



Numéro de catalogue: 11948-1-AP

Informations de base

| | | |
|--|--|--|
| Numéro de catalogue: 11948-1-AP | Numéro d'acquisition GenBank: BC008992 | Méthode de purification: Purification par affinité contre l'antigène |
| Taille: 150ul , Concentration: 450 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard; | Identification du gène (NCBI): 55816 | Dilutions recommandées: WB 1:500-1:2000 |
| Hôte: Lapin | Nom complet: docking protein 5 | |
| Isotype: IgG | MW calculé: 306 aa, 35 kDa | |
| Immunogen Catalog Number: AG2615 | MW observés: 35 kDa | |

Applications

| | |
|--|---|
| Applications testées: WB, ELISA | Contrôles positifs: WB : tissu hépatique de souris, |
| Spécificité de l'espèce: Humain, rat, souris | |

Informations générales

DOK5 belongs to the docking family of proteins that contain tandem pleckstrin homology-phosphotyrosine binding (PH-PTB) domains at their amino termini (PMID: 19909370). DOK proteins are also known as INS receptor substrates (IRS) and serve as substrates for INS and INS-like growth factor (PMID: 12730241) as well as various protein tyrosine kinases (PMID: 16647839). DOK5 functions as a specific signal transduction molecule and plays an important role in cellular differentiation. DOK5 is upregulated in systemic Sclerosis and Associated with IGFBP-5-Induced fibrosis (PMID: 24551065).

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



mouse liver tissue were subjected to SDS PAGE followed by western blot with 11948-1-AP (DOK5 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.