

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-FAM107A



Numéro de catalogue: **12176-1-AP**

Phare

4 Publications

## Informations de base

Numéro de catalogue:	BC010561	Méthode de purification:
12176-1-AP		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
150ul , Concentration: 500 µg/ml by Nanodrop and 173 µg/ml by Bradford method using BSA as the standard;	11170	IHC 1:50-1:500 IF 1:10-1:100
Hôte:	Nom complet:	
Lapin	family with sequence similarity 107, member A	
Isotype:	MW calculé	
IgG	144 aa, 17 kDa	
Immunogen Catalog Number:		
AG2820		

## Applications

Applications testées:	Contrôles positifs:
IF, IHC, ELISA	IHC : tissu de cancer du poumon humain,
Demandes citées:	IF : cellules MCF-7,
IF, IHC, WB	
Spécificité de l'espèce:	
Humain, souris	
Espèces citées:	
Humain, souris	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>	

## Informations générales

FAM107A, also named as TU3A, encodes a 144 amino-acid protein, which contains a coiled-coil domain and a nuclear localization signal. FAM107A as a transcription factor regulates some gene transcription and signal transduction in cell. With the exception of peripheral blood cells, FAM107A is widely expressed in normal tissues, but shows significant loss of expression in renal cell carcinomas and frequent loss of expression in cervical, gastric, ovarian and non small cell lung cancers. Transfection of FAM107A mRNA into cancer cell line inhibits cell growth and proliferation, supporting the evidence that the gene functions as an important tumor suppressor.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Jian Zhang	29632473	Front Mol Neurosci	WB
Bo Wan	30137324	Hum Mol Genet	WB
Shuai Ke	36010909	Cancers (Basel)	WB,IHC

## Stockage

### Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C


\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 12176-1-AP (FAM107A antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 12176-1-AP (FAM107A antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of MCF-7 cells, using FAM107A antibody 12176-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).