

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-DNAJC1



Numéro de catalogue: 12362-1-AP

Informations de base

Numéro de catalogue: 12362-1-AP	Numéro d'acquisition GenBank: BC030955	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop and 500 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 64215	Dilutions recommandées: WB 1:2000-1:10000 IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB IHC 1:50-1:500
Hôte: Lapin	Nom complet: DnaJ (Hsp40) homolog, subfamily C, member 1	
Isotype: IgG	MW calculé: 554 aa, 64 kDa	
Immunogen Catalog Number: AG3030	MW observés: 64 kDa	

Applications

Applications testées:

IHC, IP, WB, ELISA

Spécificité de l'espèce:

Humain, rat, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules MCF-7, cellules A549, cellules NIH/3T3, cellules Raji

IP : cellules Raji,

IHC : tissu de cancer du sein humain, tissu de cancer du foie humain

Informations générales

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 12362-1-AP (DNAJC1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




A549 cells were subjected to SDS PAGE followed by western blot with 12362-1-AP (DNAJC1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




IP result of anti-DNAJC1 (IP:12362-1-AP, 4ug; Detection:12362-1-AP 1:1000) with Raji cells lysate 3200 ug.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 12362-1-AP (DNAJC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 12362-1-AP (DNAJC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12362-1-AP (DNAJC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12362-1-AP (DNAJC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).