

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Alix



Numéro de catalogue: 12422-1-AP

Phare

150 Publications

Informations de base

Numéro de catalogue:	BC020066	Méthode de purification:
12422-1-AP	Identification du gène (NCBI):	Purification par affinité contre l'antigène
Taille:	10015	Dilutions recommandées:
150ul, Concentration: 800 µg/ml by Nanodrop;	Nom complet:	WB 1:5000-1:50000
Hôte:	programmed cell death 6 interacting protein	IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Lapin	MW calculé	IHC 1:50-1:500
Isotype:	868 aa, 96 kDa	IF 1:10-1:100
IgG	MW observés:	
Immunogen Catalog Number:	96 kDa, 75-80 kDa	
AG3074		

Applications

Applications testées:	Contrôles positifs:
IF, IHC, IP, WB, ELISA	WB : cellules HEK-293, cellules HeLa, cellules Jurkat, cellules NIH/3T3, tissu hépatique de rat, tissu hépatique de souris
Demandes citées:	IP : cellules Jurkat,
IF, IHC, IP, WB	IHC : tissu de cancer du foie humain,
Spécificité de l'espèce:	IF : cellules HeLa,
Humain, rat, souris	
Espèces citées:	
Humain, rat, souris	
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.	

Informations générales

ALG-2-interacting protein 1 (ALIX), also known as AIP1 or Hp95, is encoded by PDCD6IP gene and is involved in cell death through mechanisms involving its binding partner ALG-2 (apoptosis-linked gene-2). ALG-2 is a 22-kDa protein containing five serially repetitive EF-hand structures and is defined as a regulator of calcium-induced apoptosis following endoplasmic reticulum (ER) stress. ALIX interacts with ALG-2 through its C-terminal proline-rich region and participates in formation of multivesicular bodies. Recent finding suggest that ALIX is a critical component of caspase 9 activation and apoptosis triggered by calcium. The alix antibody recognizes an additional band of 75-80 kDa which has also been observed in cells and exosomes.

Publications notables

Autrice	Pubmed ID	Journal	Application
Christopher R Silvers	34564696	Br J Cancer	WB
Denghui Wei	32958903	Cell Res	WB
Xue Zou	33002418	Mol Ther	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 12422-1-AP (Alix antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



IP Result of anti-PDCD6IP (IP:12422-1-AP, 3ug; Detection:12422-1-AP 1:500) with Jurkat cells lysate 4000ug.



WB result of Alix antibody (12422-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Alix transfected HEK-293 cells.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12422-1-AP (Alix antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).