

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Cadherin-13



Numéro de catalogue: 12618-1-AP

Informations de base

Numéro de catalogue:
12618-1-AP

Taille:
150ul, Concentration: 300 µg/ml by Nanodrop and 220 µg/ml by Bradford method using BSA as the standard;

Hôte:
Lapin

Isotype:
IgG

Immunogen Catalog Number:
AG3295

Numéro d'acquisition GenBank:
BC030653

Identification du gène (NCBI):
1012

Nom complet:
cadherin 13, H-cadherin (heart)

MW calculé:
713 aa, 78 kDa

MW observés:
105 kDa

Méthode de purification:
Purification par affinité contre l'antigène

Dilutions recommandées:
WB 1:500-1:1000
IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
IHC 1:20-1:200

Applications

Applications testées:
FC, IHC, IP, WB, ELISA

Spécificité de l'espèce:
Humain, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6.0.

Contrôles positifs:

WB : tissu cardiaque de souris, cellules L02, tissu cardiaque humain

IP : tissu cardiaque de souris,

IHC : tissu de cancer de l'endomètre humain, tissu cardiaque humain

Informations générales

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. Cadherin-13 (CDH13, also known as H-cadherin or T-cadherin) is an atypical member of cadherin family, lacking transmembrane and cytosolic domains and anchored to the exterior surface of the plasma membrane via a GPI anchor. CDH13 is thought to affect cellular behavior largely through its signaling properties. It is highly expressed in heart, and in the CNS, CDH13 is expressed in cerebral cortex, medulla, hippocampus, amygdala, thalamus and substantia nigra. It is often down-regulated in cancerous cells. The CDH13 cDNA encodes a 713-amino acid protein, which has a calculated molecular weight of 78 kDa. Studies have detected variable bands on Western blots, including 105 kDa (mature form), 130 kDa (precursor), and 45 kDa, which may be a post-translational processing variant of CDH13 (PMID: 19399994).

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




mouse heart tissue were subjected to SDS PAGE followed by western blot with 12618-1-AP (Cadherin-13 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human endometrial cancer using 12618-1-AP (Cadherin-13 antibody) at dilution of 1:100 (under 10x lens).



1X10⁶ Jurkat cells were stained with 0.5ug Cadherin-13 antibody (12618-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



IP Result of anti-Cadherin-13 (IP:12618-1-AP, 4ug; Detection:12618-1-AP 1:700) with mouse heart tissue lysate 4000ug.