

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-DFNA5/GSDME



Numéro de catalogue: 13075-1-AP

Phare

22 Publications

Informations de base

Numéro de catalogue:

13075-1-AP

Taille:

150ul, Concentration: 500 µg/ml by Nanodrop;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG3746

Numéro d'acquisition GenBank:

BC019689

Identification du gène (NCBI):

1687

Nom complet:

deafness, autosomal dominant 5

MW calculé

496 aa, 55 kDa

MW observés:

55 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:2000-1:10000

IHC 1:50-1:500

Applications

Applications testées:

FC, IHC, WB, ELISA

Demandes citées:

CoIP, IF, IHC, WB

Spécificité de l'espèce:

Humain, souris

Espèces citées:

Humain, souris

Contrôles positifs:

WB : cellules A549, cellules HeLa, cellules SH-SY5Y, cellules Y79

IHC : tissu cérébral de souris, tissu d'intestin grêle de souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Informations générales

DFNA5 (deafness, autosomal dominant 5), also known as GSDME or ICERE-1, is a 496 amino acid protein that is expressed in cochlea tissue, as well as in placenta, brain, heart, liver, lung and pancreas. Defects in the gene encoding DFNA5 are the cause of non-syndromic sensorineural deafness autosomal dominant type 5 (DFNA5), a form of sensorineural hearing loss that results from damage to one of various structures that receive sound information in the brain.

Publications notables

Autrice	Pubmed ID	Journal	Application
YuanLi Huang	34594133	Cancer Manag Res	IHC
Yuan-Li Huang	34553845	Cancer Rep (Hoboken)	IHC
Xiaolin Zhong	36100190	Brain Res Bull	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 13075-1-AP (DFNA5/GSDME antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




WB result of DFNA5/ GSDME antibody (13075-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-DFNA5/ GSDME transfected A549 cells.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13075-1-AP (DFNA5 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13075-1-AP (DFNA5 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10⁶ SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human DFNA5/GSDME (13075-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).