

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-LNX1



Numéro de catalogue: 13124-1-AP

Informations de base

Numéro de catalogue: 13124-1-AP	Numéro d'acquisition GenBank: BC022983	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul , Concentration: 300 µg/ml by Nanodrop and 133 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 84708	Dilutions recommandées: WB 1:500-1:2400
Hôte: Lapin	Nom complet: ligand of numb-protein X 1	
Isotype: IgG	MW calculé: 728 aa, 81 kDa	
Immunogen Catalog Number: AG3823	MW observés: 70 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : tissu testiculaire de souris, tissu rénal de souris
Spécificité de l'espèce: Humain, rat, souris	

Informations générales

LNX1 (Ligand of Numb, protein X 1) was originally identified as a binding partner of the cell-fate determinant Numb and was reported to be involved in ubiquitylation and degradation of Numb. LNX1 contains four PDZ domains and may function as a scaffold for a multi-protein complex. LNX1 has two splicing variants, LNX1p70 and LNX1p80. LNX1p70 has the N-terminal Numb binding region and four PDZ domains, whereas LNX1p80 has an additional RING-finger domain at the N-terminus followed by a Numb-binding region and four PDZ domains. This antibody can detect both LNX1p70 and LNX1p80, around 70-80 kDa and 80-100 kDa, respectively.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



mouse testis tissue were subjected to SDS PAGE followed by western blot with 13124-1-AP (LNK1 antibody) at dilution of 1:70 incubated at room temperature for 1.5 hours.