

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-CAMKK1



Numéro de catalogue: 13624-1-AP

Informations de base

Numéro de catalogue:

13624-1-AP

Taille:

150ul, Concentration: 750 µg/ml by Nanodrop and 327 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG4534

Numéro d'acquisition GenBank:

BC031647

Identification du gène (NCBI):

84254

Nom complet:

calcium/calmodulin-dependent protein kinase kinase 1, alpha

MW calculé

520 aa, 57 kDa

MW observés:

63 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:500-1:1000

IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB

IHC 1:100-1:1000

Applications

Applications testées:

IHC, IP, WB, ELISA

Spécificité de l'espèce:

Humain, rat, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6.0.

Contrôles positifs:

WB: cellules Jurkat,

IP: cellules Jurkat,

IHC: tissu de gliome humain, tissu d'estomac humain

Informations générales

Calcium/calmodulin-dependent protein kinase kinase 1 catalyzes the phosphorylation and activation of Ca(2+)/calmodulin kinase (CaMK) as an upstream component of the CaMK cascade that has been implicated in neuronal gene transcription, synaptic plasticity, and long-term memory consolidation. CAMKK1 has two isoforms of 56kda and 58kda and involved in regulating cell apoptosis, promotes cell survival by phosphorylating AKT1/PKB that inhibits pro-apoptotic BAD/Bcl2-antagonist of cell death. Researches showed that Camkk1 plays a selective role in contextual fear memory. This 66-68kda protein can be phosphorylated (probably antophosphorylation) (PMID:7961813). This antibody recognize the N-terminal of CAMKK1.

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Jurkat cells were subjected to SDS PAGE followed by western blot with 13624-1-AP (CAMKK1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 13624-1-AP (CAMKK1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 13624-1-AP (CAMKK1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-CAMKK1 (IP:13624-1-AP, 4ug; Detection:13624-1-AP 1:1000) with Jurkat cells lysate 4000ug.