

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-SYNCRIP



Numéro de catalogue: 14024-1-AP

Phare

2 Publications

Informations de base

Numéro de catalogue:

14024-1-AP

Taille:

150ul, Concentration: 400 µg/ml by Nanodrop;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG5167

Numéro d'acquisition GenBank:

BC032643

Identification du gène (NCBI):

10492

Nom complet:

synaptotagmin binding, cytoplasmic RNA interacting protein

MW calculé

623 aa, 70 kDa

MW observés:

64-74 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:500-1:2000

IP 0.5-4.0 µg for IP and 1:500-1:1000 for WB

IF 1:50-1:500

Applications

Applications testées:

IF, IP, WB, ELISA

Demandes citées:

IP, WB

Spécificité de l'espèce:

Humain

Espèces citées:

Humain, souris

Contrôles positifs:

WB : cellules A431, cellules HEK-293, cellules HeLa, cellules Jurkat

IP : cellules A431,

IF : cellules A431,

Informations générales

Synaptotagmin-binding, cytoplasmic RNA-interacting protein(SYNCRIP), also identified as hnRNP Q, involves in mRNA processing mechanisms. It's component of the CRD-mediated complex that promotes MYC mRNA stability. Also it is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. As previously reported, the 65 and 74 kDa proteins corresponded to two isoforms of SYNCRIP produced from alternatively spliced mRNAs (PMID: 15340051).

Publications notables

Autrice	Pubmed ID	Journal	Application
Evan P Booy	30247708	Nucleic Acids Res	WB,IP
Silvia Parisi	33504840	Sci Rep	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




A431 cells were subjected to SDS PAGE followed by western blot with 14024-1-AP (SYNCRIP antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP Result of anti-SYCRIP (IP:14024-1-AP, 4ug; Detection:14024-1-AP 1:500) with A431 cells lysate 1200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 14024-1-AP (SYNCRIP antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 14024-1-AP (SYNCRIP antibody) at dilution of 1:9000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using SYCRIP antibody (14024-1-AP) at dilution of 1:200 and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). DAPI (blue).