

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-RNF114



Numéro de catalogue: 14338-1-AP

Informations de base

Numéro de catalogue: 14338-1-AP	Numéro d'acquisition GenBank: BC066919	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul , Concentration: 500 µg/ml by Nanodrop;	Identification du gène (NCBI): 55905	Dilutions recommandées: WB 1:1000-1:8000
Hôte: Lapin	Nom complet: ring finger protein 114	
Isotype: IgG	MW calculé: 26 kDa	
Immunogen Catalog Number: AG5649	MW observés: 26 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules Jurkat,
Spécificité de l'espèce: Humain	

Informations générales

RNF114, also known as ZNF228 or ZNF313, is a member of the zinc-binding protein family. RNF114 contains two C2H2 and Ring finger zinc-binding domains and a ubiquitin-interacting motif (UIM). RNF114 functions as an E3 ligase in the ubiquitin-proteasome system and plays an important role in basic cellular processes in different types of cells. RNF114 is expressed widely in many normal tissues, such as the heart, liver, and kidney. RNF114 has 2 isoforms with the molecular mass of 21 and 26 kDa. RNF114 is a novel determinant of psoriasis susceptibility, and its overexpression is a crucial indicator of epithelial inflammation (PMID: 34104941, 35069903, 37523804).

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Jurkat cells were subjected to SDS PAGE followed by western blot with 14338-1-AP (RNF114 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.