

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-ACADVL



Numéro de catalogue: 14527-1-AP

Phare

8 Publications

Informations de base

Numéro de catalogue:	BC012912	Méthode de purification:
14527-1-AP		Purification par affinité contre l'antigène
Taille:	37	Dilutions recommandées:
150ul , Concentration: 350 µg/ml by Nanodrop and 273 µg/ml by Bradford method using BSA as the standard;	Nom complet:	WB 1:1000-1:4000 IHC 1:100-1:400
Hôte:	acyl-Coenzyme A dehydrogenase, very long chain	
Lapin	MW calculé	
Isotype:	70 kDa	
IgG	MW observés:	
Immunogen Catalog Number:	66-73 kDa	
AG6016		

Applications

Applications testées:	Contrôles positifs:
IHC, WB,ELISA	WB : tissu hépatique de souris, cellules HeLa, cellules LO2, tissu hépatique de rat
Demandes citées:	IHC : tissu de cancer du foie humain,
IHC, WB	
Spécificité de l'espèce:	
Humain, rat, souris	
Espèces citées:	
Humain, rat, souris	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>	

Informations générales

Very long-chain acyl-CoA dehydrogenase (VLCAD) is one of four flavoproteins which catalyze the initial step of the mitochondrial b-oxidation spiral. It belongs to the acyl-CoA dehydrogenase family and is a homodimer of a 71-kDa polypeptide which contains 2 mol of FAD/mol of enzyme. The molecular mass of the nondenatured trypsinized VLCAD is 98 kDa, by gel filtration chromatography, indicating that it is a homodimer of the 48 kDa(tryptic digest) polypeptide(PMID:9461620). Defects in ACADVL are the cause of acyl-CoA dehydrogenase very long chain deficiency (ACADVLD). It has 2 isoforms(70 kDa and 68 kDa) produced by alternative splicing and a transit peptide. This antibody is specific to ACADVL.

Publications notables

Autrice	Pubmed ID	Journal	Application
De Huang	25242319	Cell Rep	WB
Qian Sun	27634671	Biochim Biophys Acta	WB
Yiyi Ma	36425583	Front Pharmacol	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 14527-1-AP (ACADVL antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of ACADVL antibody (14527-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACADVL transfected HeLa cells.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 14527-1-AP (ACADVL antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 14527-1-AP (ACADVL antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).