

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-HMOX2



Numéro de catalogue: 14817-1-AP

6 Publications

## Informations de base

Numéro de catalogue:  
14817-1-AP

Taille:  
150ul, Concentration: 600 µg/ml by Nanodrop and 340 µg/ml by Bradford method using BSA as the standard;

Hôte:  
Lapin

Isotype:  
IgG

Immunogen Catalog Number:  
AG6575

Numéro d'acquisition GenBank:  
BC002396

Identification du gène (NCBI):  
3163

Nom complet:  
heme oxygenase (decycling) 2

MW calculé  
36 kDa

MW observés:  
36 kDa

Méthode de purification:  
Purification par affinité contre l'antigène

Dilutions recommandées:  
WB 1:500-1:1000  
IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB  
IHC 1:50-1:500  
IF 1:50-1:500

## Applications

Applications testées:  
IF, IHC, IP, WB, ELISA

Demandes citées:  
IHC, WB

Spécificité de l'espèce:  
Humain, rat, souris

Espèces citées:  
Humain, rat, souris

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) A défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

Contrôles positifs:

WB : cellules HepG2, cellules K-562, cellules Raji, tissu splénique de souris

IP : cellules Raji,

IHC : tissu de lymphome humain,

IF : cellules HeLa,

## Informations générales

Heme oxygenase is the rate-limiting enzyme in the heme degradative pathway. There are 2 isozymic forms of heme oxygenase, an inducible heme oxygenase-1 (HMOX1) and a constitutive heme oxygenase-2 (HMOX2). Down-regulation of HMOX2 is associated with the increased expression of HMOX1 in human cell lines. HMOX2 is also named as HO-2 and could be implicated in the production of carbon monoxide in brain where it could act as a neurotransmitter.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Katarzyna Magierowska	26460608	PLoS One	WB
Dongqing Guo	26621031	J Cell Sci	WB
Katarzyna Magierowska	32408627	Cells	WB

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




HepG2 cells were subjected to SDS PAGE followed by western blot with 14817-1-AP (HMOX2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 14817-1-AP (HMOX2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 14817-1-AP (HMOX2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-HMOX2 (IP:14817-1-AP, 4ug; Detection:14817-1-AP 1:500) with Raji cells lysate 2000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 14817-1-AP (HMOX2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).