

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-UBR7



Numéro de catalogue: 16903-1-AP

Informations de base

Numéro de catalogue: 16903-1-AP	Numéro d'acquisition GenBank: BC015046	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul , Concentration: 600 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 55148	Dilutions recommandées: WB 1:2000-1:10000
Hôte: Lapin	Nom complet: ubiquitin protein ligase E3 component n-recognin 7 (putative)	
Isotype: IgG	MW calculé: 48 kDa	
Immunogen Catalog Number: AG10445	MW observés: 48-60 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules HeLa, cellules HepG2
Spécificité de l'espèce: Humain	

Informations générales

UBR7 (C14orf130), is a 425-amino-acid putative E3 ubiquitin-protein ligase which is a component of the N-end rule pathway. The N-end rule pathway is a ubiquitin-dependent system where E3 ligases called N-recognins, including UBR1, UBR2 and UBR7 (N-recognin-7), leading to target protein ubiquitination and subsequent degradation (PMID: 19008229). UBR7 contains a PHD-type zinc finger and a UBR-type zinc finger domain (PMID: 14702039), and probably undergoes acetylation (PMID: 19413330). Till date its precise function is not well known. Proteintech's UBR7 antibody 16903-1-AP is able to detect UBR7 with a theoretical MW of 48 kDa.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 16903-1-AP (UBR7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.