

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Factor XIIIa



Numéro de catalogue: 17223-1-AP

Informations de base

Numéro de catalogue: 17223-1-AP	Numéro d'acquisition GenBank: BC027963	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul, Concentration: 133 µg/ml by Nanodrop and 133 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 2162	Dilutions recommandées: WB 1:200-1:1000
Hôte: Lapin	Nom complet: coagulation factor XIII, A1 polypeptide	
Isotype: IgG	MW calculé: 732 aa, 83 kDa	
Immunogen Catalog Number: AG11059	MW observés: 80-85 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : tissu pulmonaire de souris,
Spécificité de l'espèce: Humain, rat, souris	

Informations générales

Factor XIIIa is a blood proenzyme that catalyzes intermolecular cross-linking of fibrinogen and is involved in clotting cascade. It has been identified in various tissues including placenta, uterus, and prostate. Anti-Factor XIIIa has been found to be useful in differentiating between dermatofibroma (90% positive) and desmoplastic malignant melanoma (negative). Factor XIIIa positivity is also seen in capillary hemangioblastoma, hemangioendothelioma, hemangiopericytoma, xanthogranuloma, xanthoma, hepatocellular carcinoma, glomus tumor, and meningioma.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



mouse lung tissue were subjected to SDS PAGE followed by western blot with 17223-1-AP (Factor XIIIa antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.