

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Histone H1.0



Numéro de catalogue: 17510-1-AP

11 Publications

Informations de base

Numéro de catalogue:

17510-1-AP

Taille:

150ul, Concentration: 500 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG9982

Numéro d'acquisition GenBank:

BC000145

Identification du gène (NCBI):

3005

Nom complet:

H1 histone family, member 0

MW calculé

21 kDa

MW observés:

32 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:500-1:1000

IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB

IF 1:50-1:500

Applications

Applications testées:

IF, IP, WB, ELISA

Demandes citées:

IF, WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

Humain, rat, souris

Contrôles positifs:

WB : cellules HeLa, cellules A431, cellules Jurkat, tissu splénique de rat, tissu splénique de souris

IP : cellules A431,

IF : cellules MCF-7,

Informations générales

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. Linker histones are involved in the formation of higher order structure in chromatin and the maintenance of overall chromatin compaction. The H1FO histones are found in cells that are in terminal stages of differentiation or that have low rates of cell division. Histone H1.0 (H1FO, H1FV) is a linker histone that is widely expressed in many tissues and almost all vertebrates, unlike some other linker histones. The observed molecular weight of H1FO is about 32 kDa.

Publications notables

Autrice	Pubmed ID	Journal	Application
Kohsuke Kato	34489496	Sci Rep	WB
Jianjian Zhang	34707090	Cell Death Discov	WB
Nan Tian	31762817	J Cancer	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 17510-1-AP (Histone H1.0 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Histone H1.0 antibody (17510-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP Result of anti-Histone H1.0 (IP:17510-1-AP, 4ug; Detection:17510-1-AP 1:500) with A431 cells lysate 2400ug.