

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-DRD1

Numéro de catalogue: 17934-1-AP

Phare

13 Publications



Informations de base

| | | |
|---|--------------------------------|---|
| Numéro de catalogue: | BC074978 | Méthode de purification: |
| 17934-1-AP | 1812 | Purification par affinité contre l'antigène |
| Taille: | Identification du gène (NCBI): | Dilutions recommandées: |
| 150ul , Concentration: 800 µg/ml by Nanodrop; | dopamine receptor D1 | WB 1:500-1:2000 |
| Hôte: | Nom complet: | IHC 1:50-1:500 |
| Lapin | dopamine receptor D1 | IF 1:50-1:500 |
| Isotype: | MW calculé | |
| IgG | 446 aa, 49 kDa | |
| Immunogen Catalog Number: | MW observés: | |
| AG12366 | 50-75 kDa | |

Applications

| | |
|---|---|
| Applications testées: | Contrôles positifs: |
| IF, IHC, WB, ELISA | WB : tissu cérébral de souris incubé à 37 °C, |
| Demandes citées: | IHC : tissu cérébral de souris, tissu cérébral de rat |
| IF, IHC, WB | IF : tissu cérébral de souris, |
| Spécificité de l'espèce: | |
| Humain, rat, souris | |
| Espèces citées: | |
| Humain, rat, souris | |
| Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0. | |

Informations générales

Dopamine is a neurotransmitter that plays a crucial role in physical and mental health, such as cardiovascular, hormonal, renal and central nervous systems (PMID: 29220802). Five subtypes of mammalian dopamine receptors are grouped into two classes, the D1- and D2-like classes. The D1-like class includes D1 and D5 receptors whereas the D2-like class includes D2, D3, D4 subtypes (PMID: 9457173). Dopamine receptor D1 (DRD1) is the most abundant form of dopamine receptor in the central nervous system. DRD1 stimulates adenylate cyclase, modulates D2 receptor activity, regulates neuron growth and differentiation, and mediates several behavioral responses (PMID: 1977312). DRD1 has a calculated molecular weight of 49 kDa, larger apparent molecular weight of 60-80 kDa may be due to glycosylation (PMID: 1281547; 23821371).

Publications notables

| Autrice | Pubmed ID | Journal | Application |
|------------------------|-----------|---------------|-------------|
| Soheil Kazemi Roodsari | 35625897 | Biomedicines | WB |
| Kyungri Kim | 33801790 | Genes (Basel) | WB |
| Yiqing Yan | 25594175 | Cell | WB |

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)



E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées




37°C incubated mouse brain tissue were subjected to SDS PAGE followed by western blot with 17934-1-AP (DRD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 17934-1-AP (DRD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).


Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 17934-1-AP (DRD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using DRD1 antibody (17934-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using DRD1 antibody (17934-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 17934-1-AP (DRD1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).