

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-FANCA-Specific

Numéro de catalogue: **19542-1-AP**



Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
19542-1-AP	NM_000135	Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
150ul , Concentration: 300 µg/ml by Nanodrop and 280 µg/ml by Bradford method using BSA as the standard;	2175	WB 1:200-1:1000
Hôte:	Nom complet:	
Lapin	Fanconi anemia, complementation group A	
Isotype:	MW calculé	
IgG	163 kDa	
	MW observés:	
	163 kDa	

Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules HeLa, cellules HEK-293
Spécificité de l'espèce:	
Humain	

Informations générales

FANCA, also named as FAA, FACA and FANCH, is DNA repair protein that may operate in a postreplication repair or a cell cycle checkpoint function. FANCA may be involved in interstrand DNA cross-link repair and in the maintenance of normal chromosome stability. Defects in FANCA are a cause of Fanconi anemia (FA) which is a genetically heterogeneous, autosomal recessive disorder characterized by progressive pancytopenia, a diverse assortment of congenital malformations, and a predisposition to the development of malignancies. The antibody is specific to isoform1 of FANCA.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



HeLa cells were subjected to SDS PAGE followed by western blot with 19542-1-AP (FANCA-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.