

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-Integrin Alpha 4



Numéro de catalogue: 19676-1-AP

Phare

16 Publications

## Informations de base

Numéro de catalogue:

19676-1-AP

Taille:

150ul, Concentration: 600 µg/ml by Nanodrop and 353 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Numéro d'acquisition GenBank:

NM\_000885

Identification du gène (NCBI):

3676

Nom complet:

integrin, alpha 4 (antigen CD49D, alpha 4 subunit of VLA-4 receptor)

MW calculé

115 kDa

MW observés:

140-150 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:6000

IHC 1:20-1:200

## Applications

Applications testées:

FC, IHC, WB, ELISA

Demandes citées:

IF, IHC, WB

Spécificité de l'espèce:

Humain

Espèces citées:

Humain, rat, souris

Contrôles positifs:

WB : cellules Jurkat,

IHC : tissu d'amygdalite humain,

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

## Informations générales

ITGA4, also named as CD49D, belongs to the integrin alpha chain family. ITGA4 is a receptor for fibronectin, VCAM1 and MADCAM1. ITGA4 recognizes one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. It recognizes the sequence Q-I-D-S in VCAM1. It recognizes the sequence L-D-T in MADCAM1. On activated endothelial cells ITGA4 triggers homotypic aggregation for most VLA-4-positive leukocyte cell lines. It may also participate in cytolytic T-cell interactions with target cells.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Li Wu	36184060	Vascul Pharmacol	IF
Yiyin Zhang	36384635	J Nanobiotechnology	WB
Vincent Yi-Fong Su	34074039	Int J Mol Sci	WB

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

**\*\*\* Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




Jurkat cells were subjected to SDS PAGE followed by western blot with 19676-1-AP (Integrin alpha-4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 19676-1-AP (Integrin alpha-4 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 19676-1-AP (Integrin alpha-4 Antibody) at dilution of 1:50.



$1 \times 10^6$  Jurkat cells were intracellularly stained with 0.4 ug Anti-Human Integrin Alpha 4 (19676-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).