

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-FOXA1



Numéro de catalogue: 20411-1-AP

8 Publications

## Informations de base

Numéro de catalogue:	BC033890	Méthode de purification:
20411-1-AP		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
150ul , Concentration: 500 µg/ml by Nanodrop;	3169	WB 1:1000-1:4000
Hôte:	Nom complet:	
Lapin	forkhead box A1	
Isotype:	MW calculé	
IgG	473 aa, 49 kDa	
Immunogen Catalog Number:	MW observés:	
AG14243	50 kDa	

## Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules MCF-7, cellules HepG2, cellules MDA-MB-468 et cellules T-47D
Demandes citées:	
ColP, WB	
Spécificité de l'espèce:	
Humain	
Espèces citées:	
Humain, souris	

## Informations générales

Forkhead box A1(FOXA1), also named hepatocyte nuclear factor 3-alpha (HNF-3A), is a transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). Proposed to play a role in translating the epigenetic signatures into cell type-specific enhancer-driven transcriptional programs. Its differential recruitment to chromatin is dependent on distribution of histone H3 methylated at 'Lys-5' (H3K4me2) in estrogen-regulated genes. Involved in the development of multiple endoderm-derived organ systems such as liver, pancreas, lung and prostate; FOXA1 and FOXA2 seem to have at least in part redundant roles (By similarity). Modulates the transcriptional activity of nuclear hormone receptors. Is involved in ESR1-mediated transcription; required for ESR1 binding to the NKK2-1 promoter in breast cancer cells; binds to the RPRM promoter and is required for the estrogen-induced repression of RPRM. Involved in regulation of apoptosis by inhibiting the expression of BCL2. Involved in cell cycle regulation by activating expression of CDKN1B, alone or in conjunction with BRCA1. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Zipeng Xie	32931885	Cancer Lett	WB
Lijun Li	36064451	Stem Cell Res Ther	WB
Wentao Liu	35081978	J Exp Clin Cancer Res	WB

## Stockage

### Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C


\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 20411-1-AP (FOXA1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 20411-1-AP (FOXA1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).