

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-CXCL9/MIG



Numéro de catalogue: 22355-1-AP

5 Publications

Informations de base

Numéro de catalogue:
22355-1-AP

Taille:
150ul, Concentration: 450 µg/ml by Nanodrop and 320 µg/ml by Bradford method using BSA as the standard;

Hôte:
Lapin

Isotype:
IgG

Immunogen Catalog Number:
AG17932

Numéro d'acquisition GenBank:
BC095396

Identification du gène (NCBI):
4283

Nom complet:
chemokine (C-X-C motif) ligand 9

MW calculé
14 kDa

MW observés:
14 kDa

Méthode de purification:
Purification par affinité contre l'antigène

Dilutions recommandées:
WB 1:500-1:2000

Applications

Applications testées:
WB, ELISA

Demandes citées:
IHC, WB

Spécificité de l'espèce:
Humain

Espèces citées:
Humain, souris

Contrôles positifs:

WB : IFN gamma and LPS treated THP-1 cells,

Informations générales

CXCL9 is a small cytokine known as MIG. CXCL9 is a T-cell chemoattractant, which is induced by IFN-γ. It is closely related to two other CXC chemokines called CXCL10 and CXCL11, whose genes are located near the gene for CXCL9 on human chromosome 4. A high level of CXCL9 revealed in peripheral liquids, can be considered as a marker of host immune response, especially of that involving Th1 cells.

Publications notables

Autrice	Pubmed ID	Journal	Application
Chengming Liu	34868057	Front Immunol	IHC
Wenqin Feng	35562800	Cancer Cell Int	IHC
Xinhang Dong	34110956	J Plast Surg Hand Surg	IHC

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



IFN gamma and LPS treated THP-1 cells were subjected to SDS PAGE followed by western blot with 22355-1-AP (CXCL9/MIG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.