

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-ASB5



Numéro de catalogue: 24031-1-AP

Informations de base

Numéro de catalogue: 24031-1-AP	Numéro d'acquisition GenBank: BC065710	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul, Concentration: 153 µg/ml by Nanodrop and 153 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 140458	Dilutions recommandées: WB 1:1000-1:4000
Hôte: Lapin	Nom complet: ankyrim repeat and SOCS box-containing 5	
Isotype: IgG	MW calculé: 329 aa, 36 kDa	
Immunogen Catalog Number: AG21230	MW observés: 36-40 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : tissu de muscle squelettique de souris, tissu cardiaque de souris, tissu de muscle squelettique de rat
Spécificité de l'espèce: Humain, souris	

Informations générales

ASB5, also named as Ankyrin repeat and SOCS box-containing 5, is a 329 amino acid protein, which contains a conserved one C-terminal SOCS box motif. Asb5 was significantly upregulated in growing collateral arteries on mRNA and protein level. It has been shown that asb5 is a protein implicated in the initiation of arteriogenesis.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 24031-1-AP (ASB5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 24031-1-AP (ASB5 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.