

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-SYNJ1



Numéro de catalogue: 24677-1-AP

3 Publications

## Informations de base

Numéro de catalogue:

24677-1-AP

Taille:

150ul, Concentration: 700 µg/ml by Nanodrop and 293 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG20364

Numéro d'acquisition GenBank:

BC136603

Identification du gène (NCBI):

8867

Nom complet:

synaptojanin 1

MW calculé

1612 aa, 178 kDa

MW observés:

140 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:8000

IHC 1:50-1:500

IF 1:20-1:200

## Applications

Applications testées:

IF, IHC, WB, ELISA

Demandes citées:

CoIP, IP, WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

Humain, souris

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) A défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.**

Contrôles positifs:

WB : tissu cérébral de souris, cellules NIH/3T3, tissu cérébral de rat

IHC : tissu cérébral humain, tissu cérébral de rat, tissu de muscle squelettique de rat, tissu de muscle squelettique humain

IF : cellules NIH/3T3,

## Informations générales

Synaptojanin 1, a polyphosphoinositide phosphatase, is expressed as two major alternatively spliced isoforms of 145 kDa (SJ145) and 170 kDa (SJ170), which are thought to have pleiotropic roles in endocytosis, signaling and actin function. SJ145 is highly enriched in nerve terminals where it participates in clathrin-dependent synaptic vesicle recycling. SJ170, which differs from SJ145 by the presence of a carboxy-terminal extension, is the predominant isoform in developing neurons and is expressed in a variety of tissues.(PMID:10801423).

## Publications notables

Autrice	Pubmed ID	Journal	Application
Saifeldin N Shehata	30880224	Cell Signal	WB,IP
David Furkert	32783964	Cell Chem Biol	WB
Yueqin Tian	37419331	Behav Brain Res	WB,CoIP

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




mouse brain tissue were subjected to SDS PAGE followed by western blot with 24677-1-AP (SYNJ1 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 24677-1-AP (SYNJ1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human brain tissue slide using 24677-1-AP (SYNJ1 Antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human brain tissue slide using 24677-1-AP (SYNJ1 Antibody) at dilution of 1:200 (under 40x lens).




Immunohistochemical analysis of paraffin-embedded rat skeletal muscle tissue slide using 24677-1-AP (SYNJ1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using 24677-1-AP (SYNJ1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded rat skeletal muscle tissue slide using 24677-1-AP (SYNJ1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 24677-1-AP (SYNJ1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 24677-1-AP (SYNJ1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).