

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-UHRF2



Numéro de catalogue: 25710-1-AP

1 Publications

## Informations de base

Numéro de catalogue:

25710-1-AP

Taille:

150ul, Concentration: 300 µg/ml by Nanodrop;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG22519

Numéro d'acquisition GenBank:

BC028397

Identification du gène (NCBI):

115426

Nom complet:

ubiquitin-like with PHD and ring finger domains 2

MW calculé

90 kDa

MW observés:

90 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:200-1:1000 for WB

## Applications

Applications testées:

IP, WB, ELISA

Demandes citées:

WB

Spécificité de l'espèce:

Humain

Espèces citées:

Humain

Contrôles positifs:

WB : cellules HL-60, cellules HeLa, cellules Jurkat, cellules PC-3

IP : cellules Jurkat,

## Informations générales

Ubiquitin-Like with PHD and ring finger domains 2 (UHRF2), a member that belongs to the family of UHRF, contains five recognizable functional domains, namely the ubiquitin-like domain (UBL) domain, tandem-Tudor domain (TTD), plant homeodomain (PHD), SET and RING associated (SRA) domain, and really interesting new gene (RING) finger domain. Due to the complex structure, UHRF2 possesses multiple functions in diverse cellular processes. As a ubiquitin E3 ligase, UHRF2 could ubiquitinate PCNP, a nuclear protein that contains two remarkable PEST sequences which are rich in proline (P), glutamic acid (E), serine (S), and threonine (T). It has been also reported that UHRF2 could serve as a vital cell cycle regulator by interacting with multiple cyclins, CDKs, p53, pRB and PCNA. UHRF2 has been revealed to possess epigenetic regulation function and is capable of maintaining 5mC levels in certain genomic loci in brain and stabilizes TIP60 to regulate H3K9ac and H3K14ac through RING finger domain. Moreover, UHRF2 could promote DNA damage repair by reducing the level of p21 mediated by RING finger domain. Recently, emerging evidence indicated that UHRF2 was involved in the tumorigenesis and progression of several human cancers, such as esophageal squamous cell carcinoma, lung cancer and colorectal cancer. UHRF2 has 2 isoforms with the molecular mass of 56 and 90 kDa. (PMID: 34400880)

## Publications notables

Autrice	Pubmed ID	Journal	Application
Shengjun Geng	35732617	Cell Death Dis	WB

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 25710-1-AP (UHRF2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-UHRF2 (IP:25710-1-AP, 4ug; Detection:25710-1-AP 1:300) with Jurkat cells lysate 4000ug.