

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-NF-M



Numéro de catalogue: 25805-1-AP 11 Publications

Informations de base

Numéro de catalogue:	BC002421	Méthode de purification:
25805-1-AP		Purification par affinité contre l'antigène
Taille:	4741	Dilutions recommandées:
150ul , Concentration: 260 µg/ml by Nanodrop and 213 µg/ml by Bradford method using BSA as the standard;		WB 1:2000-1:10000 IHC 1:50-1:500 IF 1:50-1:500
Hôte:	neurofilament, medium polypeptide	
Lapin	MW calculé	
Isotype:	102 kDa	
IgG	MW observés:	
	140 kDa	
Immunogen Catalog Number:		
AG22709		

Applications

Applications testées:	Contrôles positifs:
FC, IF, IHC, WB, ELISA	WB : tissu cérébral de souris, tissu cérébral de rat
Demandes citées:	IHC : tissu de cervelet de souris, tissu cérébral de souris, tissu cérébral humain, tissu de côlon humain
IF, IHC, WB	IF : tissu cérébral de souris,
Spécificité de l'espèce:	
Humain, rat, souris	
Espèces citées:	
canin, Humain, rat, souris	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>	

Informations générales

NEFM, also named as NEF3 and NFM, belongs to the intermediate filament family. Neurofilaments are the 10nm intermediate filaments found specifically in neurons. They are a major component of the cell's cytoskeleton, and provide support for normal axonal radial growth. Neurofilaments usually contain three intermediate filament proteins: L, M, and H which are involved in the maintenance of neuronal caliber. The names given to the three major neurofilament subunits are based upon the apparent molecular weight of the mammalian subunits on SDS-PAGE:NF-L, 65-68 kDa; NF-M,145-160 kDa and NF-H, 200-220 kDa. This antibody recognizes endogenous NF-M protein.

Publications notables

Autrice	Pubmed ID	Journal	Application
Natasha L Pacheco	29090078	Mol Autism	WB
Dong Sun	31642560	Cell Biol Int	WB
Jipeng Jiang	33026366	Biomater Sci	IF

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 25805-1-AP (NF-M antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 25805-1-AP (NF-M antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 25805-1-AP (NF-M antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 25805-1-AP (NF-M antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 25805-1-AP (NEFM Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1×10^6 PC-12 cells were intracellularly stained with 0.4 ug Anti-Human NF-M (25805-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).