

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-c-Fos



Numéro de catalogue: 26192-1-AP

Phare

26 Publications

Informations de base

Numéro de catalogue:	BC004490	Méthode de purification:
26192-1-AP	Identification du gène (NCBI):	Purification par affinité contre l'antigène
Taille:	2353	Dilutions recommandées:
150ul , Concentration: 247 µg/ml by Bradford method using BSA as the standard;	Nom complet:	WB 1:500-1:1000 IHC 1:50-1:500
Hôte:	MW calculé	
Lapin	41 kDa	
Isotype:	MW observés:	
IgG	65 kDa	
Immunogen Catalog Number:		
AG24340		

Applications

Applications testées:	Contrôles positifs:
IHC, WB, ELISA	WB : cellules RAW 264.7,
Demandes citées:	IHC : tissu cérébral de souris, tissu cérébral de rat
IF, IHC, WB	
Spécificité de l'espèce:	
Humain, rat, souris	
Espèces citées:	
canin, Humain, rat, souris	
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>	

Informations générales

c-Fos, also named as FOS and G0/G1 switch regulatory protein 7, is a 380 amino acid protein, which contains 1 bZIP (basic-leucine zipper) domain and belongs to the bZIP family. c-Fos is expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, c-Fos undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the c-Fos protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the c-FOS protein becomes nuclear. c-Fos is a very short-lived intracellular protein, which is very easy to degrade. The calculated molecular weight of c-Fos is 40 kDa, but Phosphorylated c-Fos protein is about 60-65 kDa. It is involved in important cellular events, including cell proliferation, differentiation and survival; genes associated with hypoxia; and angiogenesis; which makes its dysregulation an important factor for cancer development. It can also induce a loss of cell polarity and epithelial-mesenchymal transition, leading to invasive and metastatic growth in mammary epithelial cells. Expression of c-Fos is an indirect marker of neuronal activity because c-Fos is often expressed when neurons fire action potentials. Upregulation of c-Fos mRNA in a neuron indicates recent activity.

Publications notables

Autrice	Pubmed ID	Journal	Application
Kun Lu	29285195	Oncol Lett	IF
Disi Bai	30542609	Toxicol Res (Camb)	WB
Yu-Zhe Li	36442651	Neuropharmacology	IF

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 26192-1-AP (c-Fos antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 26192-1-AP (c-Fos antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.