

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-NKAP



Numéro de catalogue: 27437-1-AP

## Informations de base

Numéro de catalogue: 27437-1-AP	Numéro d'acquisition GenBank: BC015354	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul, Concentration: 270 µg/ml by Nanodrop;	Identification du gène (NCBI): 79576	Dilutions recommandées: WB 1:500-1:2000
Hôte: Lapin	Nom complet: NFKB activating protein	
Isotype: IgG	MW calculé 415 aa, 47 kDa	
Immunogen Catalog Number: AG26637	MW observés: 47-50 kDa	

## Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules A549, cellules HeLa, cellules Jurkat, cellules K-562
Spécificité de l'espèce: Humain	

## Informations générales

NKAP is a multifunctional nuclear protein that is involved in the activation of the ubiquitous transcription factor NF-kappaB. NKAP is associated with the histone deacetylase HDAC3 and with the Notch corepressor complex, and it thereby acts as a transcriptional repressor of Notch target genes. It is also required for alphabeta T cell development. A related pseudogene has been identified on chromosome X, while a related and intronless retrocopy, which has an intact CDS and may be functional, is located on chromosome 6.

## Stockage

**Stockage:**  
Stocker à -20°C. Stable pendant un an après l'expédition.  
**Tampon de stockage:**  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

**\*\*\* Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 27437-1-AP (NKAP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.