

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Alpha Taxilin



Numéro de catalogue: 27558-1-AP

Phare

1 Publications

Informations de base

Numéro de catalogue:
27558-1-AP

Taille:
150ul, Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

Hôte:
Lapin

Isotype:
IgG

Immunogen Catalog Number:
AG23001

Numéro d'acquisition GenBank:
BC103823

Identification du gène (NCBI):
200081

Nom complet:
taxilin alpha

MW calculé:
546 aa, 62 kDa

MW observés:
75 kDa

Méthode de purification:
Purification par affinité contre l'antigène

Dilutions recommandées:
WB 1:500-1:2000

Applications

Applications testées:
WB, ELISA

Demandes citées:
IF, WB

Spécificité de l'espèce:
Humain

Espèces citées:
Humain

Contrôles positifs:

WB : cellules HEK-293, cellules HeLa, cellules HepG2, cellules Jurkat

Informations générales

Alpha-taxilin is a member of the taxilin family which consist of at least three members: alpha-, beta- and gamma-taxilins. Alpha-taxilin was identified as a novel binding partner of the syntaxin family, which is involved in intracellular vesicle traffic. Alpha-taxilin is ubiquitously expressed. It plays an essential role for release of HBV-DNA containing particles. Alpha-taxilin encompasses 546 aa and has a calculated molecular weight of 62 kDa. It migrates on SDS-PAGE with an apparent molecular weight of 75 kDa. (PMID: 12558796; 23816704)

Publications notables

Autrice	Pubmed ID	Journal	Application
Dandan Ma	35119360	Elife	WB,IF

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various cells were subjected to SDS PAGE followed by western blot with 27558-1-AP (TXLNA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.