

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-LIMD1



Numéro de catalogue: 28106-1-AP

## Informations de base

Numéro de catalogue:

28106-1-AP

Taille:

150ul , Concentration: 900 µg/ml by Nanodrop and 467 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG27974

Numéro d'acquisition GenBank:

NM\_014240

Identification du gène (NCBI):

8994

Nom complet:

LIM domains containing 1

MW calculé

72 aa

MW observés:

72 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:8000

IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB

IHC 1:200-1:800

IF 1:200-1:800

## Applications

Applications testées:

IF, IHC, IP, WB, ELISA

Spécificité de l'espèce:

Humain

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (\*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

Contrôles positifs:

WB : cellules A549, cellules HeLa

IP : cellules HeLa,

IHC : tissu cérébral de souris,

IF : cellules HeLa,

## Informations générales

LIMD1 is a member of the Ajuba family of LIM domain-containing proteins, these kinds of proteins have been shown to play a role in intracellular signaling, transcriptional regulation and cellular differentiation, proliferation and migration (PMID:17174104). LIMD1 predominantly localizes to the cytoplasm especially in the E-cadherin cell-cell adhesive junction, yet also translocates to the nucleus when functions as an RB corepressor (PMID:19060205). And LIMD1 acts as a scaffolding protein to form a PHD-LIMD1-VHL axis, facilitating HIF1α ubiquitination and degradation by the proteasome (PMID:22800800). LIMD1 specifically interacts with retinoblastoma protein (pRB), inhibits E2F-mediated transcription, and suppresses the expression of the majority of genes with E2F1-responsive elements. As a tumor suppressor, LIMD1 blocks tumor growth in vitro and in vivo, and is frequently down-regulated in human lung tumors (PMID:15542589), and it also be detected in normal and breast cancer tissues. LIMD1 protein exists several phosphorylation sites, which may affect its theoretical molecular weight when tested. Proteintech's 28106-1-AP antibody was generated by 133 amino acids, it could recognize the full-length protein in applications.

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 28106-1-AP (LIMD1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-LIMD1 (IP:28106-1-AP, 4ug; Detection:28106-1-AP 1:1000) with HeLa cells lysate 2640 ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 28106-1-AP (LIMD1 antibody) at dilution of 1:400 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 28106-1-AP (LIMD1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).