

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Phospho-DLK/MAP3K12 (Thr43)



Numéro de catalogue: 28277-1-AP

Informations de base

Numéro de catalogue: 28277-1-AP	Numéro d'acquisition GenBank: NM_001193511	Méthode de purification: Purification par affinité contre l'antigène
Taille: 100ul , Concentration: 300 µg/ml by Nanodrop and 200 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 7786	Dilutions recommandées: WB 1:500-1:1000
Hôte: Lapin	Nom complet: mitogen-activated protein kinase kinase kinase 12	
Isotype: IgG	MW calculé: 93 kda	
	MW observés: 93 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules PC-3 traitées au PDGF,
Spécificité de l'espèce: Humain	

Informations générales

MAP3K12, also named as DLK ((dual leucine zipper kinase), is a part of a non-canonical MAPK signaling pathway. It has been reported that MAP3K12 can be activated by APOE through a receptor-mediated mechanism. MAP3K12 enhances the AP-1-mediated transcription of APP via a MAP kinase signal transduction pathway composed of MAP2K7, MAPK1/ERK2 and MAPK3/ERK1. MAP3K12 is required for pancreatic beta-cell proliferation during postnatal development and stress-induced JNK signaling in neurons.(PMID: 28111074, 25341110, 32189007)

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Non-treated PC-3 and PDGF (HZ-1215) treated PC-3 cells were subjected to SDS PAGE followed by western blot with 28277-1-AP (Phospho-DLK (Thr43) antibody) at dilution of 1:600 incubated at 4°C overnight.