

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TRPC4AP

Numéro de catalogue:**29310-1-AP**



Informations de base

| | | |
|---|---|---|
| Numéro de catalogue: | BC013144 | Méthode de purification: |
| 29310-1-AP | | Purification par affinité contre l'antigène |
| Taille: | Identification du gène (NCBI): | Dilutions recommandées: |
| 150ul , Concentration: 380 µg/ml by Nanodrop; | 26133 | WB 1:500-1:3000 |
| Hôte: | Nom complet: | |
| Lapin | transient receptor potential cation channel, subfamily C, member 4 associated protein | |
| Isotype: | MW calculé | |
| IgG | 797 aa, 91 kDa | |
| Immunogen Catalog Number: | MW observés: | |
| AG29860 | 85-90 kDa | |

Applications

| | |
|--------------------------|-------------------------------------|
| Applications testées: | Contrôles positifs: |
| WB, ELISA | WB : cellules HepG2, cellules HuH-7 |
| Spécificité de l'espèce: | |
| Humain | |

Informations générales

TRPC4AP, also named as C20orf188 and TRRP4AP, is a substrate-specific adapter of a DCX(DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complex required for cell cycle control. TRPC4AP participates in the activation of NFKB1 in response to ligation of TNFRSF1A, possibly by linking TNFRSF1A to the IKK signalosome. The calculated molecular weight of TRPC4AP is 91 kDa. Western blot analysis detected a protein of about 82 kDa expressed in several mouse cell lines and in a human promonocytic cell line.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 29310-1-AP (TRPC4AP antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.