

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-GPR161



Numéro de catalogue: **29328-1-AP**

Informations de base

Numéro de catalogue:	BC028163	Méthode de purification:
29328-1-AP	Identification du gène (NCBI):	Purification par affinité contre l'antigène
Taille: 150ul , Concentration: 400 µg/ml by Nanodrop;	23432	Dilutions recommandées:
Hôte: Lapin	Nom complet: G protein-coupled receptor 161	WB 1:500-1:2000 IF 1:200-1:800
Isotype: IgG	MW calculé 529 aa, 59 kDa	
Immunogen Catalog Number: AG30998	MW observés: 70 kDa	

Applications

Applications testées:	Contrôles positifs:
IF, WB, ELISA	WB : tissu cérébral de souris, tissu cérébral de rat
Spécificité de l'espèce: Humain, rat, souris	IF : ARPE-19 cells, cellules hTERT-RPE1

Informations générales

GPR161 (also known as RE2) is an orphan G protein-coupled receptor and plays an important role in the Hh pathway. In the absence of Hh signals, GPR161 localizes to primary cilia and keeps the downstream GLI transcription factors in their repressor forms (PMID: 26305592). Ciliary localization of GPR161 requires TULP3 and the IFT-A complex. In the presence of SHH, GPR161 is removed from primary cilia and is internalized into recycling endosomes, preventing its activity and allowing activation of the Shh signaling. GPR161 has a calculated molecular mass of 59 kDa. The 70-kDa band detected by this antibody probably represents a modified version of GPR161 (PMID: 18250320).

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 29328-1-AP (GPR161 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (-20°C Methanol) fixed ARPE-19 cells using GPR161 antibody (29328-1-AP) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Coralite®594 ARL13B antibody (CL594-17711, red).