

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TNFSF15

Numéro de catalogue: 29899-1-AP



Informations de base

Numéro de catalogue:	BC074941	Méthode de purification:
29899-1-AP		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
150ul , Concentration: 600 µg/ml by Nanodrop;	9966	WB 1:500-1:1000
Hôte:	Nom complet:	
Lapin	tumor necrosis factor (ligand) superfamily, member 15	
Isotype:	MW calculé	
IgG	28 kDa	
	MW observés:	
	28 kDa	

Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : tissu de côlon de souris, tissu de côlon de rat, tissu d'intestin grêle de souris
Spécificité de l'espèce:	
Humain, rat, souris	

Informations générales

Tumor necrosis factor ligand superfamily member 15 (TNFSF15) is also named as TL1 and VEGI. And it belongs to the tumor necrosis factor family. It is a receptor for TNFRSF25 and TNFRSF6B. VEGI induced degradation of IκB alpha, and nuclear translocation of p65 subunit of NF-κB by activating NF-κB (PMID: 10597252). TL1A mediates activation and apoptosis of NF-κB in DR3-expressing cell lines (PMID: 11911831). Overexpression by cancer cells of a secretable VEGI fusion protein resulted in abrogation of xenograft tumor progression. The isoforms show endothelial cell-specific expression and are generated from a 17 kb human gene by alternative splicing (PMID: 11923219). In addition, VEGI can inhibit vascular endothelial growth and angiogenesis (in vitro) (PMID: 9872942).

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 29899-1-AP (TNFSF15 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.