

À des fins de recherche uniquement

# Anticorps Monoclonal anti-BRD8

Numéro de catalogue: 60121-1-Ig **Phare**



## Informations de base

Numéro de catalogue: 60121-1-Ig	Numéro d'acquisition GenBank: BC008039	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 1467 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 10902	CloneNo.: 5B6D6
Hôte: Mouse	Nom complet: bromodomain containing 8	Dilutions recommandées: WB 1:500-1:2000 IHC 1:50-1:500 IF 1:1000-1:4000
Isotype: IgG1	MW calculé 120 kDa	
Immunogen Catalog Number: AG0790	MW observés: 120 kDa	

## Applications

Applications testées: IF, IHC, WB, ELISA	Contrôles positifs:
Spécificité de l'espèce: Humain, rat, souris	WB : tissu cérébral humain fœtal, cellules HeLa
<b>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</b>	IHC : tissu de cancer du côlon humain, IF : cellules HepG2, cellules HEK-293

## Informations générales

BRD8 is one of the bromodomain-containing proteins, which is an acetylated lysine-binding domain and thought to be involved in regulation of protein acetylation and/or HAT (histone acetyl transferase) activity [PMID: 12963728]. It is also reported to be an accessory subunit of NuA4 HAT complex (also known as TRRAP/TIP60 complex) through biochemical purification [PMID: 16049979]. In addition, it can bind to thyroid hormone receptor-β and/or retinoid X receptor, and can act as a coactivator of the nuclear hormone receptor-mediated transcription in reporter assays [PMID: 10517671].

## Stockage

Stockage:  
Stocker à -20°C. Stable pendant un an après l'expédition.  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




WB result of BRD8 antibody (60121-1-Ig, 1:8000) with si-Control and si-BRD8 transfected HeLa cells..



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 60121-1-Ig (BRD8 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 60121-1-Ig (BRD8 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using BRD8 antibody (60121-1-Ig, Clone: 5B6D6) at dilution of 1:2000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).