

À des fins de recherche uniquement

Anticorps Monoclonal anti-IL-24

Numéro de catalogue: 60139-1-Ig **1 Publications**



Informations de base

Numéro de catalogue: 60139-1-Ig	Numéro d'acquisition GenBank: BC009681	Méthode de purification: Précipitation de l'acide caprylique/du sulfate d'ammonium
Taille: 150ul , Concentration: 500 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 11009	CloneNo.: 2D6C6
Hôte: Mouse	Nom complet: interleukin 24	Dilutions recommandées: WB 1:500-1:1000
Isotype: IgG2a	MW calculé 207 aa, 24 kDa	
Immunogen Catalog Number: AG8282		

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : Protéine recombinante,
Demandes citées: WB	
Spécificité de l'espèce: Humain	
Espèces citées: Humain	

Informations générales

IL24, also named as MDA-7, MDA7 or ST16, belongs to the IL-10 family. It has antiproliferative properties on melanoma cells and may contribute to terminal cell differentiation. The induction of IL24 by oncogenes may support tumor growth at the early stages of cancer. The observed molecular weight of IL24 is reported between 25 and 35 kDa, a value higher than the deduced molecular weight of 23 kDa. This difference is likely due to post-translational modifications, such as glycosylation of the endogenously produced molecule.

Publications notables

Autrice	Pubmed ID	Journal	Application
Chunming Wang	33274495	J Clin Lab Anal	WB

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Recombinant protein were subjected to SDS PAGE followed by western blot with 60139-1-Ig (IL-24 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.