

À des fins de recherche uniquement

Anticorps Monoclonal anti-ZAP70

Numéro de catalogue: 60200-2-Ig **Phare**



Informations de base

Numéro de catalogue: 60200-2-Ig	Numéro d'acquisition GenBank: BC039039	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 447 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 7535	CloneNo.: 3B3C6
Hôte: Mouse	Nom complet: zeta-chain (TCR) associated protein kinase 70kDa	Dilutions recommandées: WB 1:500-1:2000
Isotype: IgG1	MW calculé 70 kDa	
Immunogen Catalog Number: AG7943	MW observés: 70 kDa	

Applications

Applications testées: WB,ELISA	Contrôles positifs: WB : cellules Jurkat,
Spécificité de l'espèce: Humain, souris	

Informations générales

Zeta-associated protein (ZAP)-70, a member of the Syk family of tyrosine kinases, plays an important role in T-cell receptor signaling, natural killer (NK) cell activation and early B-cell development. ZAP-70 is expressed in a subset of cases of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL) with unmutated immunoglobulin heavy-chain variable region (IgVH) genes and is associated with poor clinical outcome.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



WB result of ZAP70 antibody (60200-2-Ig, 1:5000) with si-control and si-ZAP70 transfected Jurkat cells..



Jurkat cells were subjected to SDS PAGE followed by western blot with 60200-2-Ig (ZAP70 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.