

À des fins de recherche uniquement

Anticorps Monoclonal anti-FAF1

Numéro de catalogue: 60231-1-Ig

Phare

2 Publications



Informations de base

Numéro de catalogue: 60231-1-Ig	Numéro d'acquisition GenBank: BC004970	Méthode de purification: Purification par protéine A
Taille: 150ul , Concentration: 780 µg/ml by Nanodrop and 607 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 11124	CloneNo.: 2E1D7
Hôte: Mouse	Nom complet: Fas (TNFRSF6) associated factor 1	Dilutions recommandées: WB 1:500-1:2000
Isotype: IgG2b	MW calculé: 650aa, 74 kDa	
Immunogen Catalog Number: AG0407	MW observés: 74 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules HEK-293,
Demandes citées: IF, WB	
Spécificité de l'espèce: Humain, rat, souris	
Espèces citées: rat, souris	

Informations générales

FAF1 was detected most abundant in testis, slightly less abundant in skeletal muscle and heart, followed by prostate, thymus, ovary, small intestine, and colon, but not in the peripheral blood leukocytes. The N-terminal region (amino acid 1-201) including the upstream ubiquitin homology domain of hFAF1 could bind with the death domain of Fas, which mediates programmed cell death, also called apoptosis, in a number of organ systems, notably the immune and nervous systems.

Publications notables

Autrice	Pubmed ID	Journal	Application
Huijing Xia	32990120	J Am Heart Assoc	WB
Jing Luo	34537481	Int Immunopharmacol	WB,IF

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



WB result of FAF1 antibody (60231-1-Ig, 1:5000) with si-Control and si-FAF1 transfected HEK-293 cells.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 60231-1-Ig (FAF1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.